

Asymptotic properties of equations in groups

Vitalii Roman'kov

Omsk State Dostoevskii University

(joint with Robert Gilman and Alexei Miasnikov)

Stevens Institute of Technology, Hoboken, USA; McGill
University, Montreal, Canada

Omsk, August 17, 2009

Free products

Usually the left side u of any equation $u = 1$ over any group G is element of a free product $G_X = G * F(X)$, where $X = \{x_1, \dots, x_k\}$ is considered as the set of variables.

We think that more naturally is to take free product in the variety generated by G .

So we assume that

$F(X)$ is a free group in the variety $\text{Var}(G)$, and $G_X = G * F(X)$ is a free product in this variety.

Free products

Usually the left side u of any equation $u = 1$ over any group G is element of a free product $G_X = G * F(X)$, where $X = \{x_1, \dots, x_k\}$ is considered as the set of variables.

We think that more naturally is to take free product in the variety generated by G .

So we assume that

$F(X)$ is a free group in the variety $\text{Var}(G)$, and $G_X = G * F(X)$ is a free product in this variety.

Equations

An **equation** $w = 1$ in k variables is defined by any element $w \in G_X$.

$SAT(G, k)$ – set of all equations (in k variables) satisfiable (have solutions) in G ,

$NSAT(G, k)$ – set of all equations (in k variables) not satisfiable in G .

Equations

An **equation** $w = 1$ in k variables is defined by any element $w \in G_X$.

$SAT(G, k)$ – set of all equations (in k variables) satisfiable (have solutions) in G ,

$NSAT(G, k)$ – set of all equations (in k variables) not satisfiable in G .

Equations

An **equation** $w = 1$ in k variables is defined by any element $w \in G_X$.

$SAT(G, k)$ – set of all equations (in k variables) satisfiable (have solutions) in G ,

$NSAT(G, k)$ – set of all equations (in k variables) not satisfiable in G .

Stratification

Let

T be a countable set equipped with a *size* (or length) function $s : T \rightarrow \mathbb{N}$ such that for every $n \in \mathbb{N}$ the *ball* $B_n = \{t \in T \mid s(t) \leq n\}$ is finite.

The size function s induces a volume stratification of the set T :

$$T = \bigcup_{r=0}^{\infty} B_r,$$

which gives a "direction" to infinity in T .

Stratification

Let

T be a countable set equipped with a *size* (or length) function $s : T \rightarrow \mathbb{N}$ such that for every $n \in \mathbb{N}$ the *ball* $B_n = \{t \in T \mid s(t) \leq n\}$ is finite.

The size function s induces a volume stratification of the set T :

$$T = \bigcup_{r=0}^{\infty} B_r,$$

which gives a "direction" to infinity in T .

Relative frequency

For a subset $A \subseteq T$ and a finite subset $B \subset T$ we define a relative frequency

$$d(A|B) = \frac{|A \cap B|}{|B|},$$

Now, one can define the r -frequency (or r -density) of A with respect to the stratification T (or the size function s) by

$$d_r(A) = d(A|B_r).$$

Relative frequency

For a subset $A \subseteq T$ and a finite subset $B \subset T$ we define a relative frequency

$$d(A|B) = \frac{|A \cap B|}{|B|},$$

Now, one can define the r -frequency (or r -density) of A with respect to the stratification T (or the size function s) by

$$d_r(A) = d(A|B_r).$$

Asymptotic density

Now, the **asymptotic density** of A with respect to the stratification T is defined as the following limit

$$ad(A) = \limsup_{r \rightarrow \infty} d_r(A)$$

If the actual limit

$$sad(A) = \lim_{r \rightarrow \infty} d_r(A)$$

exists then we call it the **strict asymptotic density** of A .

A is called **generic** if $sad(A) = 1$ and it is **negligible** if $sad(A) = 0$.

Asymptotic density

Now, the **asymptotic density** of A with respect to the stratification T is defined as the following limit

$$ad(A) = \limsup_{r \rightarrow \infty} d_r(A)$$

If the actual limit

$$sad(A) = \lim_{r \rightarrow \infty} d_r(A)$$

exists then we call it the **strict asymptotic density** of A .

A is called **generic** if $sad(A) = 1$ and it is **negligible** if $sad(A) = 0$.

Uniform asymptotic density of power sets in free abelian groups

The asymptotic density of any power set $\gamma\mathbb{Z}^k) \in \mathbb{Z}^k$ is almost obvious. But we need in estimates on the convergence rates that we could not find in the literature.

Proposition 1. Let $\gamma, k \in \mathbb{N}^+$. Then

- 1) $sad(\gamma\mathbb{Z}^k) = 1/\gamma^k$;
- 2) $|d_r(\gamma\mathbb{Z}^k) - 1/\gamma^k| \leq \frac{2^{k+1}k}{r\gamma^{k-1}}$ for every $r \geq \gamma$,
- 3) $d_r(\gamma\mathbb{Z}^k)$ converges to $1/\gamma^k$ uniformly in γ .

Primitive and γ -primitive elements of free abelian groups

An element $x = x_1^{\gamma_1} \dots x_k^{\gamma_k} \in A(X)$, where $A(X)$ is the free abelian group with basis X is called

primitive (visuable)

if and only if it is a member of some basis of $A(X)$, or, equivalently, $\gcd(\gamma_1, \dots, \gamma_k) = 1$.

It is called

γ -primitive (γ -visuable)

if and only if it is γ -power of some primitive element, or, equivalently, $\gcd(\gamma_1, \dots, \gamma_k) = \gamma$.

Asymptotic density of sets of γ -primitive elements in free abelian groups

Let $P_{k,\gamma}$ be the set of all γ -primitive elements in the free abelian group $A(X)$ of rank k .

The following result is well-known in number theory. In the case $k = 2$ it was proved by F. Mertens (1874), in full generality it is due to Christopher. Below $\zeta(k) = \sum_{n=1}^{\infty} 1/n^k$ denotes Riemann zeta-function.

Proposition 2. For each $\gamma \in \mathbf{N}$ we have

$$\text{sad}(P_{k,\gamma}) = \frac{1}{\gamma^k \zeta(k)}.$$

Uniform asymptotic density of γ -primitive sets in free abelian groups

Also we need in estimates on the convergence rates for the sets $P_{k,\gamma}$.

Proposition 3. Let $\gamma, k \in \mathbb{N}^+, \gamma \geq 2$. Then

1) For every $\varepsilon \geq 0$ there exists $r(\varepsilon) \in \mathbb{N}^+$ such that

$$|d_r(P_{k,\gamma}) - \frac{1}{\gamma^k \zeta(k)}| \leq \frac{\varepsilon}{\gamma^{k-1}}$$

for every $r \geq r(\varepsilon)$.

2) $d_r(P_{k,\gamma})$ converges to $\frac{1}{\gamma^k \zeta(k)}$ uniformly in γ .

Equations

Let

$$A = \mathbf{Z}^m$$

be a free abelian group with a basis $\{a_1, \dots, a_m\}$ ($m \geq 1$).

Now

$$X = \mathbf{Z}^k$$

is the free abelian group with a basis $\{x_1, \dots, x_k\}$ ($k \geq 1$),

and

$$A_X = A \times X = \mathbf{Z}^{m+k}$$

is the free abelian group with a basis $\{x_1, \dots, x_k, a_1, \dots, a_m\}$.

Satisfiable equations

Every element $w \in A_X$ can be uniquely written in the form

$$w = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m},$$

where $\gamma_1, \dots, \gamma_k, \alpha_1, \dots, \alpha_m \in \mathbf{Z}$.

We call $\gamma = \gcd(\gamma_1, \dots, \gamma_k)$ the **exponent** of w and denote it as $\gamma = \exp(w)$. In the exceptional case $\gamma_1 = \dots = \gamma_k = 0$ we define $\exp(w) = 0$.

Lemma 1. An equation $w = 1$ of non-zero exponent $\gamma = \exp(w)$ has a solution in A if and only if $\gamma | \gcd(\alpha_1, \dots, \alpha_m)$. For $k = 1$ and $\gamma_1 = \pm\gamma \neq 0$ there is the unique solution $x_1 = a_1^{-\alpha_1/\gamma_1} \dots a_m^{-\alpha_m/\gamma_1}$. When $\exp(w) = 0$ a solution exists if and only if $\alpha_1 = \dots = \alpha_m = 0$ (every tuple of k elements is a solution).

Satisfiable equations

Every element $w \in A_X$ can be uniquely written in the form

$$w = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m},$$

where $\gamma_1, \dots, \gamma_k, \alpha_1, \dots, \alpha_m \in \mathbf{Z}$.

We call $\gamma = \gcd(\gamma_1, \dots, \gamma_k)$ the **exponent** of w and denote it as $\gamma = \exp(w)$. In the exceptional case $\gamma_1 = \dots = \gamma_k = 0$ we define $\exp(w) = 0$.

Lemma 1. An equation $w = 1$ of non-zero exponent $\gamma = \exp(w)$ has a solution in A if and only if $\gamma | \gcd(\alpha_1, \dots, \alpha_m)$. For $k = 1$ and $\gamma_1 = \pm\gamma \neq 0$ there is the unique solution $x_1 = a_1^{-\alpha_1/\gamma_1} \dots a_m^{-\alpha_m/\gamma_1}$. When $\exp(w) = 0$ a solution exists if and only if $\alpha_1 = \dots = \alpha_m = 0$ (every tuple of k elements is a solution).

Stratification

For a free abelian group \mathbf{Z}^q a length function $l : \mathbf{Z}^q \rightarrow \mathbf{N}$ will usually be the restriction to \mathbf{Z}^q of $\|\cdot\|_\infty$ -norm from \mathbf{R}^q .

The norm $\|\cdot\|$ of an element u is defined as

$$\|u\| = \max\{|\gamma_1|, \dots, |\gamma_k|, |\alpha_1|, \dots, |\alpha_m|\}.$$

The function $l : A_X \rightarrow \mathbf{N}$ is defined as $l(u) = \|u\|$.

There are **the boxes** $B_r = \{w \in A_X : l(w) \leq r\}$, and their **slices** $B_r(\gamma) = \{w \in A_X : l(w) \leq r, \exp(w) = \gamma\}$, $\gamma = 0, 1, 2, \dots$

One-variable equations

Theorem 1. For $r, m \in \mathbb{N}^+$

$$\left| d_r(\text{SAT}(A, 1)) - \frac{Z_r(m)}{r} \right| = O\left(\frac{Z_r(m-1)}{r^2}\right).$$

Corollary 1.

The set $\text{SAT}(A, 1)$ is negligible, and $\text{NSAT}(A, 1)$ is generic.

One-variable equations

Theorem 1. For $r, m \in \mathbb{N}^+$

$$\left| d_r(\text{SAT}(A, 1)) - \frac{Z_r(m)}{r} \right| = O\left(\frac{Z_r(m-1)}{r^2}\right).$$

Corollary 1.

The set $\text{SAT}(A, 1)$ is negligible, and $\text{NSAT}(A, 1)$ is generic.

Multi-variable equations

Theorem 2. Assume that $k \geq 2, m \geq 1$. Then the set $SAT(A, k)$ has the asymptotic density

$$sad(SAT(A, k)) = \frac{\zeta(k+m)}{\zeta(k)}.$$

Basic commutators

Let $N = N_{mc}$ be the free nilpotent group of rank m and class c . Then every element $u \in N_X$ can be uniquely written in the form:

$$u = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m} \prod_{j=1}^p b_j^{\delta_j},$$

where $b_1 < \dots < b_p$ denote the set of all basic commutators. We assume that the ordering of all basic commutators of weight $j \geq 2$ is such that first s_{j-1} ones depend in a_i only, and other $p_{j-1} - s_{j-1}$ of them occur at least one of x_j .

Norm

The norm $\|\cdot\|$ of an element $u \in N_X$ is defined as

$$\|u\| = \max\{|\gamma_i|, |\alpha_l|, |\delta_j| \ (i = 1, \dots, k; l = 1, \dots, m; j = 1, \dots, p)\}.$$

The function $l : N_X \rightarrow \mathbb{N}$ is defined as $l(u) = \|u\|$. There are the **boxes**: $B_r = \{u \in N_X : l(u) \leq r\}$, and the **slices**:

$$B_{r,\gamma} = \{u \in N_X : l(u) \leq r, \gamma = \exp(u) = \gcd(\gamma_1, \dots, \gamma_k) \text{ (or } 0 \text{ if } \gamma_1 = \dots = \gamma_k = 0)\}.$$

Norm

The norm $\|\cdot\|$ of an element $u \in N_X$ is defined as

$$\|u\| = \max\{|\gamma_i|, |\alpha_l|, |\delta_j| \mid (i = 1, \dots, k; l = 1, \dots, m; j = 1, \dots, p)\}.$$

The function $l : N_X \rightarrow \mathbb{N}$ is defined as $l(u) = \|u\|$. There are the

boxes: $B_r = \{u \in N_X : l(u) \leq r\}$, and the **slices:**

$B_{r,\gamma} = \{u \in N_X : l(u) \leq r, \gamma = \exp(u) = \gcd(\gamma_1, \dots, \gamma_k) \text{ (or 0 if } \gamma_1 = \dots = \gamma_k = 0)\}$.

Main theorem

Now we can formulate our main assertions for nilpotent case.

Theorem 3. Assume that $k, m \geq 2, c \geq 2$. Then the set $SAT(N, k)$ has the asymptotic density

$$ad(SAT(N, k)) \geq \frac{\zeta(k + m + s)}{\zeta(k)}, \quad (1)$$

where s denote the total number of all basic commutators at a_1, \dots, a_m of weights $2, \dots, c - 1$.

Preliminaries

Let $F = F_m$ be a free group of rank $m \geq 2$ with basis $\mathcal{F} = \mathcal{F}_m = \{f_1, \dots, f_m\}$, and $F(X) = F_k$ be a free group of rank $k \geq 1$ with basis $X = \{x_1, \dots, x_k\}$. Then $F_X = F * F(X) = F_{m+k}$ is a space of all equations with variables from X and constants from F .

As before, F_X has the ball and spherical stratifications:

$\bigcup_{r=0}^{\infty} S_r = F_X$, $\bigcup_{r=0}^{\infty} S_r = F_X$, relative to basis $\mathcal{F} \cup X$.

Preliminaries

Let $F = F_m$ be a free group of rank $m \geq 2$ with basis $\mathcal{F} = \mathcal{F}_m = \{f_1, \dots, f_m\}$, and $F(X) = F_k$ be a free group of rank $k \geq 1$ with basis $X = \{x_1, \dots, x_k\}$. Then $F_X = F * F(X) = F_{m+k}$ is a space of all equations with variables from X and constants from F .

As before, F_X has the ball and spherical stratifications:

$$\bigcup_{r=0}^{\infty} S_r = F_X, \quad \bigcup_{r=0}^{\infty} S_r = F_X, \quad \text{relative to basis } \mathcal{F} \cup X.$$

Connection between solvability of equations in free and free abelian groups

We need to recall two known results that relate asymptotics in F_q and A_q .

Theorem by Sharp. Let $a \in A_q$ and $r \in \mathbb{N}$. Then

$$\lim_{r \rightarrow \infty} \left| \sigma^q r^{q/2} \left(\frac{|S_r(a)|}{|S_r|} + \frac{|S_{r+1}(a)|}{|S_{r+1}|} \right) - \frac{2}{(2\pi)^{q/2}} e^{-\|a\|_2^2 / 2\sigma^2 r} \right| = 0,$$

uniformly in $a \in A$.

$$\text{Here } \sigma^2 = \frac{1}{\sqrt{2q-1}} \left(1 + \left(\frac{q + \sqrt{2q-1}}{q - \sqrt{2q-1}} \right)^{1/2} \right).$$

Corollary

Corollary 1. There is a constant $c \in \mathbb{N}$ such that for any $a \in A_q$ and $r \in \mathbb{N}$

$$\frac{|S_{2r+\delta_a}(a)|}{|S_{2r+\delta_a}|} \leq \frac{c}{r^{q/2}},$$

where $\delta_a = 0$ if $\|a\|_1$ is even, and $\delta_a = 1$ if $\|a\|_1$ is odd.

Rivin's theorem

Theorem by Rivin For any $D \subseteq \mathbb{R}^q, q \geq 2$,

$$\lim_{r \rightarrow \infty} \frac{1}{|S_r|} |\{w \in S_r \mid \mu(w)/r^{1/2} \in D\}| = \frac{1}{(2\pi)^{q/2} \sigma^q} \int_D e^{-\|t\|_2^2 / 2\sigma^2} dt.$$

Asymptotic of one-variable equations

Theorem 4.

The set $SAT(F, 1)$ is negligible relative to both ball and spherical stratifications, so $sad(SAT(F, 1)) = 0$, $sad(NSAT(F, 1)) = 1$.

Split equations

We say that an equation $u = 1$, $u \in F_X$, **splits** if $u = vg^{-1}$, and so it is equivalent to equation

$$v = g,$$

where $v = v(x_1, \dots, x_k) \in F(X)$ and $g \in F$.

Denote by $V(F, k)$ the set of all split equations in k variables over F . Also let $SAT_V(F, k)$ and $NSAT_V(F, k)$ be the sets of all satisfiable and all unsatisfiable split equations from $V(F, k)$.

Conditions of satisfiability

As usual, $A_X = A \times A(X)$ is the free abelian group of rank $m + k$, the standard epimorphic image for $\mu : F_X \rightarrow F_X/F'_X = A_X$. A basis of A_X is taken as $\{a_1, \dots, a_m\}$, and $\mu(f_i) = a_i, \mu(x_j) = x_j$.

The image of an element $u \in F_X$ under μ can be uniquely written as

$$u^\mu = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m}.$$

We define $\exp(u) = \exp(u^\mu) = \gcd(\gamma_1, \dots, \gamma_k)$.

Lemma 2. Let $u \in V(F, k)$. If $\exp(u) = 1$ then $u \in \text{SAT}_V(F, k)$.

Conditions of satisfiability

As usual, $A_X = A \times A(X)$ is the free abelian group of rank $m + k$, the standard epimorphic image for $\mu : F_X \rightarrow F_X/F'_X = A_X$. A basis of A_X is taken as $\{a_1, \dots, a_m\}$, and $\mu(f_i) = a_i, \mu(x_j) = x_j$.

The image of an element $u \in F_X$ under μ can be uniquely written as

$$u^\mu = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m}.$$

We define $\exp(u) = \exp(u^\mu) = \gcd(\gamma_1, \dots, \gamma_k)$.

Lemma 2. Let $u \in V(F, k)$. If $\exp(u) = 1$ then $u \in \text{SAT}_V(F, k)$.

Conditions of satisfiability

As usual, $A_X = A \times A(X)$ is the free abelian group of rank $m + k$, the standard epimorphic image for $\mu : F_X \rightarrow F_X/F'_X = A_X$. A basis of A_X is taken as $\{a_1, \dots, a_m\}$, and $\mu(f_i) = a_i, \mu(x_j) = x_j$.

The image of an element $u \in F_X$ under μ can be uniquely written as

$$u^\mu = x_1^{\gamma_1} \dots x_k^{\gamma_k} a_1^{\alpha_1} \dots a_m^{\alpha_m}.$$

We define $\exp(u) = \exp(u^\mu) = \gcd(\gamma_1, \dots, \gamma_k)$.

Lemma 2. Let $u \in V(F, k)$. If $\exp(u) = 1$ then $u \in \text{SAT}_V(F, k)$.

Some known results

In the paper by I.Kapovich, I.Rivin, P.Schupp, V.Shpilrain, [*Densities in free groups and \mathbb{Z}^k , visible points and test elements*, Math. Research Letters, 14 (2007), no. 2, 263-284] the authors relate two densities.

They prove that if $E \subseteq \mathbb{Z}^q$ is invariant under the natural action of $SL(q, \mathbb{Z})$ then the asymptotic density of $SAT(A, k)$ in A_q and, the so called, the annular density of its full preimage $\mu^{-1}(SAT(A, k))$ in F_q are equal.

Main theorem

Assume that $k \geq 2$ and $k \geq m$. Then the asymptotic density of the set $SAT_V(F, k)$ can be estimated as follows:

Theorem 5. $ad(SAT_V(F, k)) \geq \frac{2}{(2k-1)\zeta(k)}$.

The set $NSAT(F, k)$ can be estimated too.

Main theorem

Assume that $k \geq 2$ and $k \geq m$. Then the asymptotic density of the set $SAT_V(F, k)$ can be estimated as follows:

Theorem 5. $ad(SAT_V(F, k)) \geq \frac{2}{(2k-1)\zeta(k)}$.

The set $NSAT(F, k)$ can be estimated too.